

Predicting wound complications following plastic surgeon closure of spine surgeries

E. Hope Weissler BA, Christian Piña BA, Michael Ingargiola MD, Nachi Gupta MD PhD, Felipe Molina-Burbano BS, Peter J. Taub MD

Introduction: Spine surgeries continue to grow in popularity, recently increasing by more than 200% in a decade.¹ At our institution, plastic surgeons frequently assist spine surgeons with wound closures of index spine procedures. In anticipation of this becoming a more common practice nationwide, the authors sought to determine risk factors for wound complications in this setting.

Methods: Spine surgeries closed by a single plastic surgeon at a large academic hospital were reviewed. Patients 18 years or younger, with invasiveness indices of zero,² current wound infections, or undergoing surgery for management of complications from prior procedures were excluded. Factors significantly associated ($p < 0.05$) with wound complications on univariate analysis were included in a regression model.

Results: Seven hundred eight procedures were done. Twenty-one patients had any wound complication, including 2 superficial infections, 5 deep infections, 3 dehiscences, 4 seromas, and 7 hematomas. Patients undergoing cervical surgery were less likely to have a wound complication (OR 0.39, 0.14-1.09). Patients requiring intra-operative blood transfusion (OR 3.42, 1.29-9.08) and with ASA ≥ 3 (OR 4.68, 1.70-12.92) were more likely to have a wound complication. Surgical time was longer (266 ± 141 versus 196 ± 97 minutes, $p = 0.009$) and estimated blood loss higher (1063 ± 1032 versus 615 ± 786 mL, $p = 0.021$) among patients suffering wound complications. Invasiveness index was not associated with wound complications (0.711). In a multivariate logistic regression controlling for EBL, operative time, ASA status, and intra-operative transfusion, only ASA status of 3 or greater predicted complications ($p = 0.005$). Cervical surgeries were associated with fewer wound complications on multivariate analysis (OR 0.29, 0.09-0.92).

Conclusions: Contrary to papers in the spine literature that have found operative duration, diabetes, hypertension, and age, among other risk factors, predictive of complications, we found that only ASA status of ≥ 3 was associated.^{3,4,5} This may reflect a lack of power, as the ASA classification aggregates comorbidities and BMI. Patients at increased risk for complications should be managed more aggressively, including prophylactic local muscle flap closure where appropriate.

¹ Belatti DA, Phisitkul P. Trends in orthopedics: An analysis of Medicare claims, 2000-2010. *Orthopedics*. 2013;36: e366-72.

² Mirza MK, Deyo RA, Heagerty PJ, Turner JA, Lee LA, Goodkin R. Towards standardized measurement of adverse events in spine surgery: conceptual model and pilot evaluation. *BMC Musculoskeletal Disorders*. 2006;7: doi:10.1186/1471-2474-7-53.

³ Kimmel KT, Algattas H, Joynt P et al. Risk modeling predicts complication rates for spinal surgery. *Spine*. 2015;40: 1836-1841.

⁴ Bernatz JT, Anderson PA. Thirty-day readmission rates in spine surgery: systematic review and meta-analysis. *Neurosurg Focus*. 2015;39(4): 1-9.

⁵ Bekelis K, Desai A, Bakhoum SF, Missios S. A predictive model of complications after spine surgery: The National Surgical Quality Improvement Program (NSQIP): 2005-2010. *The Spine Journal*. 2014;14: 1247-1255.